# WF4313/6413-Fisheries Management















### Carnivores in a Changing World: New Perspectives

Research Seminar 3:00pm, November 29, 2018 A100 – Tully

Claudia Wultsch, PhD City University of New York, NY, US



# Revised Schedule\*\*

- November 27<sup>th</sup> –No lab work on your briefs
- DRAFT Due 11/30/2018 by 5pm!
- December 4<sup>th</sup> Brief presentations during lab.
- Final Exam 12/11 @ 8am




# Methods of Eradication & Control


- Chemicals
  - Rotenone, Lampricide
- Physical
  - Traps, nets, explosives, water level, electrofishing, commercial fishing
- Biological
  - Predator & prey, pathogens, daughterless technologies, pheromones

# Climate change & water


- Amount
- Temperature




# Precipitation trends




# Temperature trends






# Pacific decadal oscillation






# Pacific decadal oscillation



# Salmon

- Columbia stocks
- Passing Bonneville Dam



# **UW TODAY**

January 14, 2013

### Salmon runs boom, go bust over centuries

#### Sandra Hines

News and Information

Salmon runs are notoriously variable: strong one year, and weak the next. New research shows that the same may be true from one century to the next.

Scientists in the past 20 years have recognized that salmon stocks vary not only year to year, but also on decades-long time cycles. One example is the 30-year to 80-year booms and busts in salmon runs in Alaska and on the West Coast driven by the climate pattern known as the Pacific Decadal Oscillation.

Now work led by University of Washington researchers reveals those decadal cycles may overlay even more important, centuries-long conditions, or regimes, that influence fish productivity. Cycles lasting up to 200 years were found while examining 500-year records of salmon abundance in Southwest Alaska. Natural variations in the abundance of spawning salmon are as large those due to human harvest.





Researchers gathered sediment cores from lakes in 16 major watersheds in southwestern Alaska. Lauren Rogers/U of Washington

Cycles lasting up to 200 years were found while examining 500-year records of salmon abundance in Southwest Alaska. Natural variations in the abundance of spawning salmon are as large those due to human harvest.

# A take home message

### Recruitment camps:

- 1. Climate driven
- 2. Spawner driven

Need both...

Minimize excessive recruit overfishing so when conditions are good recruitment can happen...



### Effects on fish & fisheries

- 1. Fishery closures
- 2. Movement of fish stocks
- 3. Disease
- 4. And more...

### Dozens of sturgeon found dead in Columbia River

Originally published July 16, 2015 at 6:56 am | Updated July 16, 2015 at 12:23 pm





Sophia Murillo of Kennewick, left, and Frank Carr of Sequim wade into the Columbia River Wednesday to check out an estimated seven-foot long dead sturgeon at the east end of Pasco's Wade Park near Road 39. State fish... (Bob Brawdy/Tri-City Herald) More >

Washington Fish and Wildlife officials have received repeated reports of dead sturgeon this week on the Columbia River, but the exact cause of death remains a mystery.

### Hoot owl closures

#### Heat wave spells early trouble for Montana trout

JUNE 24, 2015 BY LAURA LUNDQUIST

River levels are dropping rapidly as summer temperatures continue to climb, and both trends spell trouble for Montana's trout. As Montana Fish, Wildlife & Parks prepares for fishing closures, anglers can do some things to help fish out.

By Saturday, high temperatures are predicted to top 100 degrees and stay that way at least through Wednesday for parts of western Montana. That's 20 to 25 degrees above normal for this time of year and the extended heat wave will break records,

### **Hoot owl**

#### FISHING

#### Go fishing: Hoot owl restrictions lifted on some western Montana rivers



AUGUST 19, 2015 8:52 PM · BY PERRY BACKUS

Longer nights and cooler water temperatures mean an end to fishing restrictions on three of western Montana's most favored trout rivers.

Montana Fish, Wildlife and Parks lifted "hoot owl" restrictions on the Bitterroot, the entire main stem of the Blackfoot and the Clark Fork, downstream of the Rock Creek confluence Wednesday.

The rivers had been under the restrictions that prohibit fishing from 2 p.m. to midnight since July 3 in an effort to reduce the impact on drought-stressed trout.

FWP's Bitterroot fisheries biologist Chris Clancy said this has been an unusual summer temperature-wise.

"On most years, you could bet money and win that peak water temperatures would appear sometime in the last week of July or first week of August," Clancy said. "This year, the temperatures peaked in late June, early July. It looked really grim for trout back then."

Some timely rains and periodic breaks in the heat made a difference this summer for the Bitterroot River.

"It really didn't turn out all that badly this year," Clancy said.

The state begins to consider instituting "hoot owl" restrictions when water temperatures go over 72 degrees for three consecutive days. Temperatures need to drop below 70 degrees for the same amount of time before the restrictions are withdrawn.

While the upper reaches of the Bitterroot remained cooler due to releases from Painted Rock Reservoir this summer, Clancy said the river downstream of Hamilton started hitting 73 degrees in late June.

In the same time period, water temperatures rose to 76 degrees in the lower reaches of the Bitterroot River near Missoula.

"Trout really don't like 70-degree water all that well," Clancy said.

The preferred water temperature for rainbow and brown trout is around 55 to 65 degrees. Temperatures in the mid 60s are stressful to native bull and westslope cutthroat trout.

### Movement

#### Warmer waters shake up Shore fishing

JANUARY 4, 2016, 10:16 PM LAST UPDATED SUADAY, JANUARY 4, 2016, 10:20 PM BY JAMES M. O'NELL STAFF WINTER IT HE RECORD



PATRICK PLANGAMENTIAL TO THE RECO-Malin Pinels, a marine biologist with Rutgers University, has studied how rising ocean temperatures have affected the

commercial and regreational fishing industries of the Jersey Shore.

Species like shad have shifted to the north

A dramatic rise in ocean temperature is changing the kinds of fish that swim off New Jersey's coast = a development that has significant ramifications for the state's fishing industry and could even alter restaurant menus.

Species that used to be centered off New Jersey, like shad, have shifted north to New England in search of cooler waters. Fish once centered off Virginia, like black sea bass and summer flounder, now swim in abundance off New Jersey.

"We've been seeing unprecedented warming in the ocean over the past several decades," said Malin Pinsky, a marine biologist at Rutgers University who has studied the changes. "Climate change and the warming of ocean waters is not some abstract global problem. We can see the implications right off our shore in New Jersey."


Temperatures had risen one degree every 14 years off New Jersey since the 1970s. In the past decade, warming has accelerated to one degree every 25 years, Pinsky said.

"Marine fish are very sensitive to a change in temperature = they can only survive in a narrow range, so they are seeking out cooler waters toward the poles and deeper in the ocean," he said. "And deeper generally means farther from shore." Malin Pinsky, a marine biologist with Rutgers University, has studied how rising ocean temperatures have affected the commercial and recreational fishing industries off the Jersey Shore.


Black sea bass seen at Gray's Reef National Islanine Sancribury in Georgia. The fish once centered of the coast of Virginia are now found in abundance near


# Within stream networks

Move towards cooler water, if available...




# **Brook trout**





# Effect on fisherman



# Transport thermal barriers






### Truckin': Salmon Take A Long, Strange Trip To The Pacific Ocean

MARCH 26, 2014 3:21 AM ET





Comment

SHARE

Pacific Or Bust: Fingerling Chinook salmon are dumped into a holding pen Tuesday as they are transferred from a truck into the Sacramento River in Rio Vista, Calif. From here, they'll be towed downstream for a bit, then make their own way out

# Disease


#### Klamath River agreement

An agreement to resolve distribution of water, restore habitat and remove four hydroelectric dams on the upper Klamath River to boost the watershed's salmon population is in peril.















# Willamette basin spring Chinook


# Anadromous species of conservation need

- -Threatened status 1999
- Anthropogenic modifications







# Limited natural reproduction
















www.nwd-wc.usace.army.mil


# **Natural production**




# **Natural production**



# **Natural production**



# Reintroduction above dams




### **Problem: Prespawn mortality (PSM)**

In excess of 90% in some years

—Temperature




# **Prespawning mortality (PSM)**




### Holding in cool pathogen free water









### PSM and thermal exposure




## PSM and thermal exposure




# Race against time



# Race against time




## Race against time



## **Example management alternatives**

#### Hypothesized to reduce PSM

- 1.Trap → outplant: prioritize brood stock
- 2.Trap  $\rightarrow$  outplant: proportional allocation
- 3. Trap, hold @ hatchery → outplant





### How do we decide?











### A decision framework

### Structured decision making approach

- Simulation model
- 2. Decision model



### How do we decide?



### What are decision models?

# Focus on decision

- 1. Optimal decision
- 2. Account for <u>uncertainty</u>
- 3. Sensitivity of decision
- 4. Inform Research & Monitoring

### An example

What is the optimal decision?

